

Contents of Volume 3

COMMENTARIES

The price of lost freedom: entropy of bimolecular complex formation A.V.Finkelstein and J.Janin 1

De novo design of an α/β barrel protein

T.Handel 233

REVIEW

Catalytic antibodies—a new direction in enzyme design M.J.Powell and D.E.Hansen 69

PROTOCOLS

Rapid and efficient site-specific mutagenesis M.Baldrich and W.Goebel 563

Versatile protein engineering vectors for mutagenesis, expression and hybrid enzyme formation F.W.Larimer, R.J.Mural and T.S.Soper 227

ABSTRACTS

Protein Engineering '89 279

PROTEIN STRUCTURE AND ANALYSIS

Nuclear magnetic resonance studies of isolated structural domains of yeast phosphoglycerate kinase W.J.Fairbrother, P.Minard, L.Hall, J.-M.Betton, D.Missiakas, J.M.Yon and R.J.P.Williams 5

The structure of interfaces between subunits of dimeric and tetrameric proteins S.Miller 77

Molecular cartography of proteins: surface relief analysis of the calf eye lens protein gamma-crystallin Yu.Chirgadze, N.Kurochkina and S.Nikonov 105

Thermitase and proteinase K: a comparison of the refined three-dimensional structures of the native enzymes Ch.Betzel, A.V.Teplyakov, E.H.Harutyunyan, W.Saenger and K.S.Wilson 161

Crystallization and structure solution at 4 Å resolution of the recombinant synthase domain of N-(5'-phosphoribosyl)anthranilate isomerase:indole-3-glycerol-phosphate synthase from *Escherichia coli* complexed to a substrate analogue M.Wilmanns, E.Schlagenhauf, B.Fol and J.N.Jansonius 173

An 8-fold $\beta\alpha$ barrel protein with redundant folding possibilities K.Luger, H.Szadkowski and K.Kirschner 249

The differences in conformation between α -human atrial natriuretic polypeptide, α -hANP, and its derivative, Met(O)- α -hANP, in solution S.Koyama, Y.Kobayashi, T.Ohkubo, Y.Kyogoku, A.Sato, M.Kobayashi and N.Go 393

An attempt to unify the structure of polymerases M.Delarue, O.Poch, N.Tordo, D.Moras and P.Argos 461

Modeling the intact form of the α_1 -proteinase inhibitor R.A.Engh, H.T.Wright and R.Huber 469

Sequence requirements for cytochromes P450IIA1 and P450IIA2 catalytic activity: evidence for both specific and non-specific substrate binding interactions through use of chimeric cDNAs and cDNA expression N.Hanioka, K.Korzekwa and F.J.Gonzalez 571

Structural effects induced by removal of a disulfide-bridge: the X-ray structure of the C30A/C51A mutant of basic pancreatic trypsin inhibitor at 1.6 Å C.Eigenbrot, M.Randal and A.A.Kossiakoff 591

| | | |
|--|---|------------|
| Incorporation of crystallographic temperature factors in the statistical analysis of protein tertiary structures | R.Bott and J.Frane | 649 |
| PREDICTION OF PROTEIN STRUCTURE | | |
| A predicted three-dimensional structure for the human immunodeficiency virus binding domains of CD4 antigen | P.A.Bates, M.J.McGregor, S.A.Islam, Q.J.Sattentau and M.J.E.Sternberg | 13 |
| Amino acid sequence templates derived from recurrent turn motifs in proteins: critical evaluation of their predictive power | M.J.Rooman, S.J.Wodak and J.M.Thornton | 23 |
| A prediction of tertiary structures of peptide by the Monte Carlo simulated annealing method | H.Kawai, T.Kikuchi and Y.Okamoto | 85 |
| Stereochemical modeling of disulfide bridges. Criteria for introduction into proteins by site-directed mutagenesis | R.Sowdhamini, N.Srinivasan, B.Shoichet, D.V.Santi, C.Ramakrishnan and P.Balaram | 95 |
| Construction of validated, non-redundant composite protein sequence databases | A.J.Bleasby and J.C.Wootton | 153 |
| An object-oriented database for protein structure analysis | P.M.D.Gray, N.W.Paton, G.J.L.Kemp and J.E.Fothergill | 235 |
| A sequence motif in the transmembrane region of growth factor receptors with tyrosine kinase activity mediates dimerization | M.J.E.Sternberg and W.J.Gullick | 245 |
| Molecular modelling study of antigen binding to oxazolone-specific antibodies: the $\text{\O}1$ idotypic IgG and its mature variant with increased affinity to 2-phenyloxazolone | L.Holm, L.Laaksonen, M.Kaartinen, T.T.Teeri and J.K.C.Knowles | 403 |
| Flagellin as an object for supramolecular engineering | O.V.Fedorov and A.V.Efimov | 411 |
| Comparison of dielectric response models for simulating electrostatic effects in proteins | E.L.Mehler | 415 |
| Free energy changes associated with amino acid substitution in proteins | M.A.S.Saqi and J.M.Goodfellow | 419 |
| Protein fragment variability analysis and some principles of protein engineering of vaccines | A.M.Eroshkin, V.I.Fomin, P.A.Zhilkin and V.A.Kulichkov | 425 |
| Sequence differences between glycosylated and non-glycosylated Asn-X-Thr/Ser acceptor sites: implications for protein engineering | Y.Gavel and G.von Heijne | 433 |
| β-Turns and their distortions: a proposed new nomenclature | C.M.Wilmot and J.M.Thornton | 479 |
| Influence of secondary structure on the hydration of serine, threonine and tyrosine residues in proteins | N.Thanki, J.M.Thornton and J.M.Goodfellow | 495 |
| Energetic approach to the folding of four α-helices connected sequentially | L.Carlacci and K.-C.Chou | 509 |
| Determination of reliable regions in protein sequence alignments | M.Vingron and P.Argos | 565 |
| Alignment of protein sequences using secondary structure: a modified dynamic programming method | F.Fischel-Ghodsian, G.Mathiowitz and T.F.Smith | 577 |
| Testing the procedure of simulated annealing by refining homologous immunoglobulin light-chain dimers | Z.-B.Xu, C.-H.Chang and M.Schiffer | 583 |

| | | |
|--|--|------------|
| Predicting surface exposure of amino acids from protein sequence | S.R.Holbrook, S.M.Muskal and S.-H.Kim | 659 |
| Prediction of the disulfide-bonding state of cysteine in proteins | S.M.Muskal, S.R.Holbrook and S.-H.Kim | 667 |
| The interpretation of site-directed mutagenesis experiments by linear free energy relations | J.E.Straub and M.Karplus | 673 |
| Molecular dynamics simulations of active site mutants of triosephosphate isomerase | V.Daggett and P.A.Kollman | 677 |
| PROTEIN ENGINEERING—BY SITE-DIRECTED MUTAGENESIS | | |
| Site-directed alteration of Glu197 and Glu66 in a pyruvoyl-dependent histidine decarboxylase | H.E.McElroy and J.D.Robertus | 43 |
| Tissue-type plasminogen activator variants with domain duplications and rearrangements | W.Markland, D.Pollock and D.J.Livingston | 111 |
| Structure–function analysis of tissue-type plasminogen activator by linker-insertion, point and deletion mutagenesis | W.Markland, D.Pollock and D.J.Livingston | 117 |
| Alteration of aspartate 101 in the active site of <i>Escherichia coli</i> alkaline phosphatase enhances the catalytic activity | A.Chaidaroglou and E.R.Kantrowitz | 127 |
| Site-directed mutagenesis of <i>Escherichia coli</i> translation initiation factor IF1. Identification of the amino acids involved in its ribosomal binding and recycling | C.O.Gualerzi, R.Spurio, A.La Teana, R.Calogero, B.Celano and C.L.Pon | 133 |
| Catalytic mechanism of fungal glucoamylase as defined by mutagenesis of Asp176, Glu179 and Glu180 in the enzyme from <i>Aspergillus awamori</i> | M.R.Sierks, C.Ford, P.J.Reilly and B.Svensson | 193 |
| Introduction of internal cysteines as conformational probes in yeast phosphoglycerate kinase | N.Ballery, P.Minard, M.Desmadril, J.-M.Betton, D.Perahia, L.Mouawad, L.Hall and J.M.Yon | 199 |
| Metallothioneins with interdomain hinges expanded by insertion mutagenesis | I.-K.Rhee, K.S.Lee and P.C.Huang | 205 |
| A thermoresistant mutant of ribonuclease T1 having three disulfide bonds | S.Nishikawa, J.Adiwinata, H.Morioka, T.Fujimura, T.Tanaka, S.Uesugi, T.Hakoshima, K.Tomita, S.Nakagawa and M.Ikehara | 443 |
| Site-directed mutagenesis of aspartic acid 372 at the ATP binding site of yeast phosphoglycerate kinase: over-expression and characterization of the mutant enzyme | P.Minard, D.J.Bowen, L.Hall, J.A.Littlechild and H.C.Watson | 515 |
| Protein engineering of alcohol dehydrogenases; effects of amino acid changes at positions 93 and 48 of yeast ADH1 | E.H.Creaser, C.Murali and K.A.Britt | 523 |
| Effect on inhibitory activity of mutation at reaction site P4 of the <i>Streptomyces subtilisin</i> inhibitor, SSI | S.Kojima, I.Kumagai and K.Miura | 527 |
| Structural definition by antibody engineering of an idiotypic determinant | M.Sollazzo, D.Castiglia, R.Billetta, A.Tramontano and M.Zanetti | 531 |
| Influence of size and polarity of residue 31 in porcine pancreatic phospholipase A₂ on catalytic properties | O.P.Kuipers, J.Kerver, J.van Meersbergen, R.Vis, R.Dijkman, H.M.Verheij and G.H.de Haas | 599 |
| Protein engineering of chymosin; modification of the optimum pH of enzyme catalysis | D.Mantafounis and J.Pitts | 605 |

| | | |
|--|---|------------|
| Increased biological activity of a recombinant factor IX variant carrying alanine at position +1 | P.Meulien, A.Balland, P.Lepage, F.Mischler, K.Dott, C.Hauss, M.Grandgeorge and J.-P.Lecocq | 629 |
| Site-specific attachment to recombinant antibodies via introduced surface cysteine residues | A.Lyons, D.J.King, R.J.Owens, G.T.Yarranton, A.Millican, N.R.Whittle and J.R.Adair | 703 |
| Invariant amino acid replacement affects the dihydrofolate reductase function and its gene expression | N.V.Murzina and A.T.Gudkov | 709 |
| Mutational analysis of structure-activity relationships in human tumor necrosis factor-alpha | J.Yamagishi, H.Kawashima, N.Matsuo, M.Ohue, M.Yamayoshi, T.Fukui, H.Kotani, R.Furuta, K.Nakano and M.Yamada | 713 |
| Alteration in folding efficiency and conformation of recombinant human tumor necrosis factor-alpha by replacing cysteines 69 and 101 with aspartic acid 69 and arginine 101 | T.Arakawa, J.V.Visger, M.McGinley, M.F.Rohde, G.M.Fox and L.O.Narhi | 721 |
| PROTEIN ENGINEERING—BY RANDOM MUTAGENESIS | | |
| Random mutagenesis used to probe the structure and function of <i>Bacillus stearothermophilus</i> alpha-amylase | L.Holm, A.K.Koivula, P.M.Lehtovaara, A.Hemminki and J.K.C.Knowles | 181 |
| Mutational analysis of the C-terminus of human interferon-γ | R.Wetzel, L.J.Perry, C.Veilleux and G.Chang | 611 |
| PROTEIN ENGINEERING—BY CHEMICAL METHODS | | |
| Chemical synthesis in protein engineering: total synthesis, purification and covalent structural characterization of a mitogenic protein, human transforming growth factor-alpha | D.D.L.Woo, I.Clark-Lewis, B.T.Chait and S.B.H.Kent | 29 |
| Lipid modification of proteins and their membrane transport | A.V.Kabanov, A.V.Levashov and V.Yu.Alakhov | 39 |
| Sequential protection-modification method for selective sulphydryl group derivatization in proteins having more than one cysteine | A.Planas and J.F.Kirsch | 625 |
| CLONING AND EXPRESSION | | |
| Efficient expression and characterization of isolated structural domains of yeast phosphoglycerate kinase generated by site-directed mutagenesis | P.Minard, L.Hall, J.-M.Betton, D.Missiakas and J.M.Yon | 55 |
| Recombinant rabbit uteroglobin expressed at high levels in <i>E.coli</i> forms stable dimers and binds progesterone | W.Peter, M.Beato and G.Suske | 61 |
| Direct expression in <i>E.coli</i> of a functionally active protein A—snake toxin fusion protein | F.Ducancel, J.-C.Boulain, O.Trémeau and A.Ménéz | 139 |
| The location of inhibitory specificities in human mucus proteinase inhibitor (MPI): separate expression of the COOH-terminal domain yields an active inhibitor of three different proteinases | B.Meckelein, T.Nikiforov, A.Clemen and H.Appelhans | 215 |
| Efficient expression and Zn(II)-dependent structure of the DNA binding domain of the yeast GAL4 protein | Y.Serikawa, M.Shirakawa, H.Matsuo and Y.Kyogoku | 267 |
| Construction of rat aldolase C expression plasmid and the hybrid formation between rat aldolase C and human aldolase A or B co-expressed in <i>Escherichia coli</i> | Y.Takasaki and K.Hori | 273 |
| Expression of goat α-lactalbumin in <i>Escherichia coli</i> and its refolding to biologically active protein | I.Kumagai, S.Takeda, T.Hibino and K.Miura | 449 |

| | | |
|---|--|-----|
| Functional expression <i>in vitro</i> of bovine visual rhodopsin | S.A.Zozulya, V.V.Gurevich, T.A.Zvyaga, E.P.Shirokova, I.L.Dumler, M.N.Garnovskaya, M.Yu.Natochin, B.E.Shmukler and P.R.Badalov | 453 |
| Mammalian cell transient expression of tissue factor for the production of antigen | L.R.Paborsky, B.M.Fendly, K.L.Fisher, R.M.Lawn, B.J.Marks, G.McCray, K.M.Tate, G.A.Vehar and C.M.Gorman | 547 |
| A dual-affinity gene fusion system to express small recombinant proteins in a soluble form: expression and characterization of protein A deletion mutants | B.Jansson, C.Palmcrantz, M.Uhlén and B.Nilsson | 555 |
| Expression, purification and crystallization of penicillin G acylase from <i>Escherichia coli</i> ATCC 11195 | P.D.Hunt, S.P.Tolley, R.J.Ward, C.P.Hill and G.G.Dodson | 635 |
| Expression of mouse immunoglobulin light and heavy chain variable regions in <i>Escherichia coli</i> and reconstitution of antigen-binding activity | H.Field, G.T.Yarranton and A.R.Rees | 641 |
| Expression of human asparagine synthetase in <i>Saccharomyces cerevisiae</i> | G.Van Heeke and S.M.Schuster | 739 |
| PHYSICO-CHEMICAL STUDIES OF PROTEIN STRUCTURE AND FUNCTION | | |
| Structure and activity of artificial mutant variants of human growth hormone | S.Nishikawa, Y.Nishida, H.Uemura, Y.Yamada, T.Tanaka, S.Uesugi, M.Morikawa, E.Uchida, T.Hayakawa and M.Ikehara | 49 |
| Energetics of RNA cleavage: implications for the mechanism of action of ribozymes | K.Taira, M.Uebayasi, H.Maeda and K.Furukawa | 691 |
| SYNTHETIC GENES AND PROTEINS | | |
| Biosynthesis of winter flounder antifreeze proprotein in <i>E.coli</i> | I.D.Peters, C.L.Hew and P.L.Davies | 145 |
| A method for construction of long randomized open reading frames and polypeptides | W.Mandecki | 221 |
| Synthesis, purification and initial structural characterization of octarellin, a <i>de novo</i> polypeptide modelled on the α/β-barrel proteins | K.Goraj, A.Renard and J.A.Martial | 259 |
| Synthesis and expression in <i>Escherichia coli</i> of DNA encoding the murine λ_1 chain of a monoclonal antibody specific for <i>Salmonella</i> serotype B O-antigen | N.N.Anand, G.Dubuc, S.Mandal, J.Phipps, M.A.J.Gidney, B.Sinnott, N.M.Young, C.R.MacKenzie, D.R.Bundle and S.A.Narang | 541 |
| Improvement of nutritional value and functional properties of soybean glycinin by protein engineering | C.S.Kim, S.Kamiya, T.Sato, S.Utsumi and M.Kito | 725 |
| Construction of a novel artificial-ribosome-releasing plasmid | K.Taira, M.Oda, H.Shinshi, H.Maeda and K.Furukawa | 733 |